arXiv:1608.01655v1 [math.NT] 4 Aug 2016

ON GAUSS PERIODS

XIANG-DONG HOU

Abstract. Let q be a prime power, and let $r = nk + 1$ be a prime such that $r \nmid q$, where n and k are positive integers. Under a simple condition on q, r and k, a Gauss period of type (n, k) is a normal element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$; the complexity of the resulting normal basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is denoted by $C(n, k; q)$. Recent works determined $C(n, k; q)$ for $k \le 7$ and all qualified n and q. In this paper, we show that for any given $k > 0$, $C(n, k; q)$ is given by an explicit formula except for finitely many primes $r = nk+1$ and the exceptional primes are easily determined. Moreover, we describe an algorithm that allows one to compute $C(n, k; q)$ for the exceptional primes $r = nk + 1$. The numerical results of the paper cover $C(n, k; q)$ for $k \le 20$ and all qualified n and q.

1. Introduction and Preliminaries

We briefly recall the existing results on Gauss periods, some of which are quite recent.

Let q be a prime power, and let $r = nk + 1$ be a prime such that $r \nmid q$, where n and k are positive integers. Since $k \mid r - 1$, there is a unique cyclic subgroup K of $\mathbb{Z}_r^*$ with $|K| = k$. Since $q^{nk} - 1 = q^r - 1 \equiv 0 \pmod r$, there exists $\gamma \in \mathbb{F}_{q^{nk}}^*$ such that $o(\gamma) = r$. Let

$$\alpha_i = \sum_{a \in K} \gamma^{q^i a}, \quad 0 \le i \le n - 1, \tag{1.1}$$

which are called *Gauss periods* of type (n, k) over $\mathbb{F}_q$. Note that $\alpha_i = \alpha_0^{q^i}$. Since $(q^n)^k \equiv 1 \pmod r$, we have $q^n \in K$. Therefore

$$\alpha_i^{q^n} = \sum_{a \in K} \gamma^{q^i q^n a} = \sum_{a \in K} \gamma^{q^i a} = \alpha_i,$$

and hence $\alpha_i \in \mathbb{F}_{q^n}$.

A necessary condition for $\alpha_0, \ldots, \alpha_{n-1}$ to form a normal basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is that these elements be distinct. This condition requires $q^i K$, $0 \le i \le n - 1$, to be distinct cosets of K in $\mathbb{Z}_r^*$, i.e., $\langle q, K\rangle = \mathbb{Z}_r^*$, which happens if and only if $\gcd(nk/e, n) = 1$, where $e = e(q, r)$ is the order of q in $\mathbb{Z}_r^*$.

On the other hand, if $\langle q, K\rangle = \mathbb{Z}_r^*$, then $\alpha_0, \ldots, \alpha_{n-1}$ indeed form a normal basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$; the reason is the following [4, 13]. Assume that $\sum_{i=0}^{n-1} c_i\alpha_i = 0$, where $c_i \in \mathbb{F}_q$, i.e.,

$$\sum_{i=0}^{n-1} \sum_{a \in q^i K} c_i \gamma^a = 0.$$

Key words and phrases. cyclotomic number, Gauss period, finite field, normal basis.

Treat each $a \in \mathbb{Z}_r^*$ as an interger in $\{1, \dots, r-1\}$, and let

$$f(X) = \sum_{i=0}^{n-1} \sum_{a \in q^i K} c_i X^a \in \mathbb{F}_q[X].$$

For each $m \in \mathbb{Z}_r^*$, we have $m = q^j a_0$ for some $0 \le j \le n-1$ and $a_0 \in K$. Thus

$$f(\gamma^m) = \sum_{i=0}^{n-1} \sum_{a \in q^i K} c_i \gamma^{q^j a_0 a} = \Big(\sum_{i=0}^{n-1} \sum_{a \in q^i K} c_i \gamma^a \Big)^{q^j} = f(\gamma)^{q^j} = 0.$$

Also note that $f(0) = 0$. Therefore $f(x) = 0$ for all $x \in \{0\} \cup \langle \gamma \rangle$. Since $\deg f \le r-1$, we must have $f = 0$, i.e., $c_i = 0$ for all $0 \le i \le n-1$.

From now one, we assume that $\langle q, K \rangle = \mathbb{Z}_r^*$ and hence $\alpha_0, \dots, \alpha_{n-1}$ form a normal basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$. To avoid triviality, we also assume that $n > 1$, so nk is even. For $0 \le i \le n-1$,

$$\begin{aligned} \alpha_0 \alpha_i &= \sum_{a,b \in K} \gamma^{a+q^i b} = \sum_{a,b \in K} \gamma^{a(1+q^i b)} \\ &= k|\{-q^{-i}\} \cap K| + \sum_{j=0}^{n-1} \alpha_j |(1+q^i K) \cap q^j K|. \end{aligned} \tag{1.2}$$

In the above,

$$t_{ij} = |(1+q^i K) \cap q^j K|, \quad 0 \le i, j \le n-1, \tag{1.3}$$

are called the *cyclotomic numbers*. Note that

$$q^{-i} \in -K \Leftrightarrow \begin{cases} q^{-i} \in K \Leftrightarrow i = 0 & \text{if } k \text{ is even,} \\ q^{-2i} \in K \text{ but } q^{-i} \notin K \Leftrightarrow i = n/2 & \text{if } k \text{ is odd.} \end{cases}$$

Hence $|\{-q^{-i}\} \cap K| = \delta_i$, where

$$\delta_i = \begin{cases} 1 & \text{if } k \text{ is even and } i = 0, \text{ or } k \text{ is odd and } i = n/2, \\ 0 & \text{otherwise.} \end{cases} \tag{1.4}$$

Now (1.2) becomes

$$\alpha_0 \alpha_i = k\delta_i + \sum_{j=0}^{n-1} t_{ij} \alpha_j = \sum_{j=0}^{n-1} (t_{ij} - k\delta_i) \alpha_j, \quad 0 \le i \le n-1. \tag{1.5}$$

(The last step in the above follows from the fact that $\sum_{j=0}^{n-1} \alpha_j = \sum_{a \in \mathbb{Z}_r^*} \gamma^a = -1$.) Let $p = \operatorname{char} \mathbb{F}_q$ and let

$$C(n,k;q) = \big|\{(i,j) : 0 \le i, j \le n-1,\ t_{ij} - k\delta_i \not\equiv 0 \pmod p\}\big|, \tag{1.6}$$

which is the number of nonero entries of the matrix $(t_{ij} - k\delta_i)_{0 \le i,j \le n-1}$ over $\mathbb{F}_p$ and is called the *complexity* of the normal basis $\alpha_0, \dots, \alpha_{n-1}$ of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$. Normal bases with low complexity are sought as they provide effective algorithms for multiplication and exponentiation in finite fields [5].

Equation (1.6) involves two primes: $p = \operatorname{char} \mathbb{F}_q$ and $r = nk+1$. The prime p only plays a superficial role since the cyclotomic numbers t_{ij} depend only on r and k.

The values of $C(n,k;q)$ were first determined in [10] for $k = 1$ and for $k = 2$ with an even q. (The case $k = 2$ with an arbitrary q is covered by a recent result of [8].)

A more careful investigation of the cyclotomic numbers t_{ij} in [3] produced explicit formulas for $C(n,k;q)$ with $3 \le k \le 6$ and $n \ge 3$. More recently, the following general result was proved in [8]:

Theorem 1.1 ([8, Theorem 1.7]). *Assume that $t_{ij} \le 2$ for all $0 \le i,j \le n-1$.*

(i) *If k is even,*

$$C(n,k;q) = \begin{cases} nk - k^2 + 3k - 3 & \text{if } p = 2, \\ nk - \frac{1}{2}k^2 + \frac{3}{2}k - 2 & \text{if } p > 2 \text{ and } k \equiv 0 \pmod p, \\ n(k+1) - \frac{1}{2}k^2 + k - 3 & \text{if } p > 2 \text{ and } k \equiv 1 \pmod p, \\ n(k+1) - \frac{1}{2}k^2 + \frac{1}{2}k - 1 & \text{if } p > 2 \text{ and } k \equiv 2 \pmod p, \\ n(k+1) - \frac{1}{2}k^2 + k - 2 & \text{if } p > 2 \text{ and } k \not\equiv 0,1,2 \pmod p. \end{cases}$$

(ii) *If k is odd,*

$$C(n,k;q) = \begin{cases} n(k+1) - k^2 + k - 1 & \text{if } p = 2, \\ nk - \frac{1}{2}k^2 + \frac{3}{2}k - 2 & \text{if } p > 2 \text{ and } k \equiv 0 \pmod p, \\ n(k+1) - \frac{1}{2}k^2 - \frac{1}{2}k & \text{if } p > 2 \text{ and } k \equiv 1 \pmod p, \\ n(k+1) - \frac{1}{2}k^2 + \frac{1}{2}k - 1 & \text{if } p > 2 \text{ and } k \not\equiv 0,1 \pmod p. \end{cases}$$

It is claimed in [8] that for $k = 7$, the only exceptional case not covered by Theorem 1.1 is $n = 4$ ($r = 29$). We will see that for $k = 7$, [8] missed another exceptional case ($n = 6$, $r = 43$); for $k = 6$, [3] also missed an exceptional case ($n = 3$, $r = 19$).

The main contribution of the present paper is the following: For each given $k \ge 1$, we show that except for finitely many primes r, which we shall call *exceptional primes*, the condition $t_{ij} \le 2$ ($0 \le i,j \le n-1$) is satisfied and hence $C(n,k;q)$ is given by Theorem 1.1. The exceptional primes are easily determined using resultants in characteristic 0. Moreover, for each exceptional prime r, we describe an algorithm for computing the value distribution of the cyclotomic numbers, which then gives $C(n,k;q)$. The combined message is that for any given k (not too big), $C(n,k;q)$ can be determined for all qualified n and q. We demonstrate the computational results of $C(n,k;q)$ for $k \le 20$.

Section 2 contains a review and a further discussion of the cyclotomic numbers. In Section 3 we prove that for each given k, there are only finitely many exceptional primes r, and we explain where these exceptional primes come from and how to find them. In Section 4 we describe an algorithm that allows one to compute $C(n,k;q)$ for a given k, an exceptional prime $r = nk+1$, and all qualified q. The computational results of $C(n,k;q)$ for $k \le 20$ are included. We also give a theoretic explanation for the computational results with $n = 2$.

2. Cyclotomic Numbers

Cyclotomic numbers have been studied in terms of Jacobi sums by many authors; see for example [1, 2, 6, 7, 11, 12, 14]. The focus of the present paper is a little different. Most (but not all) of the results gathered in this section have appeared, explicitly or implicitly, in [3, 8].

Recall that $n > 1$, $r = nk + 1$ is a prime, K is the unique subgroup of $\mathbb{Z}_r^*$ of order k, and $\langle q, K\rangle = \mathbb{Z}_r^*$. Write $K_i = q^i K$, $0 \le i \le n-1$, hence

$$t_{ij} = |(1 + K_i) \cap K_j|, \quad 0 \le i, j \le n-1. \tag{2.1}$$

Lemma 2.1. *$t_{ij} > 0$ if and only if there exists $x \in \mathbb{Z}_r \setminus \{0, -1\}$ such that $K_i = xK$ and $K_j = (1 + x)K$.*

Proof. ($\Leftarrow$) We have $1 + x \in (1 + K_i) \cap K_j$, hence $t_{ij} > 0$.

($\Rightarrow$) Let $y \in (1 + K_i) \cap K_j$ and $x = y - 1$. Then $x \in K_i$ and $1 + x \in K_j$. □

For $x, y \in \mathbb{Z}_r \setminus \{0. - 1\}$, define $x \sim y$ if and only if $xK = yK$ and $(1 + x)K = (1 + y)K$, i.e., $x/y \in K$ and $(1 + x)/(1 + y) \in K$. Let $[x]$ denote the $\sim$ equivalence class of x. For conveninence, we also define $[0] = \{0\}$ and $[-1] = \{-1\}$.

Lemma 2.2. *For each $x \in \mathbb{Z}_r$,*

$$(1 + xK) \cap (1 + x)K = 1 + [x]. \tag{2.2}$$

Proof. Let $y \in \mathbb{Z}_r$. If $x \ne 0, -1$,

$$\begin{aligned} 1 + y \in (1 + xK) \cap (1 + x)K &\Leftrightarrow y \in xK \text{ and } 1 + y \in (1 + x)K \\ &\Leftrightarrow y \sim x. \end{aligned}$$

If $x = 0$,

$$1 + y \in (1 + 0K) \cap (1 + 0)K \Leftrightarrow y = 0.$$

If $x = -1$,

$$1 + y \in (1 - K) \cap 0K \Leftrightarrow y = -1.$$

□

Let $K = \langle\omega\rangle$ and let $\mathcal{S}_k = \{(u, v) \in \mathbb{Z}_k^2 : u \ne 0,\ v \ne 0,\ u \ne v\}$. Define

$$\begin{aligned} S: \quad \mathcal{S}_k &\longrightarrow \mathbb{Z}_r^* \setminus \{-1\} \\ (u, v) &\longmapsto -\frac{1 - \omega^v}{\omega^u - \omega^v}. \end{aligned} \tag{2.3}$$

Lemma 2.3. *Let $x, y \in \mathbb{Z}_r^* \setminus \{-1\}$. Then $x \sim y$ but $x \ne y$ if and only if there exists $(u, v) \in S^{-1}(x)$ such that $y = S(-u, -v)$.*

Proof. ($\Rightarrow$) Since $x \sim y$, we have

$$\begin{cases} \dfrac{y}{x} = \omega^u, \\ \dfrac{1 + y}{1 + x} = \omega^v, \end{cases} \tag{2.4}$$

for some $u, v \in \mathbb{Z}_k$. Since $x, y \in \{0, -1\}$ and $x \ne y$, we have $u \ne 0$, $v \ne 0$ and $u \ne v$. Solving (2.4) gives $x = S(u, v)$ and $y = S(-u, -v)$.

($\Rightarrow$) (2.4) is satisfied for $x = S(u, v)$ and $y = S(-u, -v)$. Since $u \ne 0$, $y \ne x$. □

Proposition 2.4. *Let $x \in \mathbb{Z}_r^* \setminus \{-1\}$. Then*

$$|[x]| = 1 + |S^{-1}(x)|. \tag{2.5}$$

Proof. By Lemma 2.3,

$$[x] = \{x\} \dot{\cup} \{S(-u,-v) : (u,v) \in S^{-1}(x)\},$$

where $\dot{\cup}$ means disjoint union. It remains to show that the mapping

$$\begin{aligned} S^{-1}(x) &\longrightarrow \mathbb{Z}_r^* \setminus \{-1\} \\ (u,v) &\longmapsto S(-u,-v) \end{aligned}$$

is one-to-one. Let $(u_1,v_1),(u_2,v_2) \in S^{-1}(x)$ be such that $S(-u_1,-v_1) = S(-u_2,-v_2)$. Then

$$\omega^{u_1} S(u_1,v_1) = S(-u_1,-v_1) = S(-u_2,-v_2) = \omega^{u_2} S(u_2,v_2).$$

It follows that $u_1 = u_2$. Then $S(u_1,v_1) = S(u_1,v_2)$, which gives $v_1 = v_2$. □

Remark 2.5. It follows from Lemma 2.1, (2.2) and (2.5) that $t_{ij} \le 2$ for all $0 \le i,j \le n-1$ if and only if the mapping $S : \mathcal{S}_k \to \mathbb{Z}_r^* \setminus \{-1\}$ is one-to-one.

For $0 \le i \le n-1$ and $0 \le \tau \le k$, define

$$a_i(\tau) = |\{0 \le j \le n-1 : t_{ij} = \tau\}|, \tag{2.6}$$

$$a(\tau) = \sum_{i=0}^{n-1} a_i(\tau). \tag{2.7}$$

Then

$$\sum_{\tau} \tau a_i(\tau) = \sum_j t_{ij} = \sum_j |(1+K_i) \cap K_j| = |(1+K_i) \cap \mathbb{Z}_r^*| = k - \delta_i \tag{2.8}$$

and

$$\sum_{\tau} \tau a(\tau) = nk - 1. \tag{2.9}$$

Let

$$a_*(\tau) = \begin{cases} a_0(\tau) & \text{if } k \text{ is even,} \\ a_{n/2}(\tau) & \text{if } k \text{ is odd.} \end{cases} \tag{2.10}$$

It follows from (1.6) that

$$\begin{aligned} C(n,k;q) &= \sum_{\tau \not\equiv 0 \,(\mathrm{mod}\, p)} \big(a(\tau) - a_*(\tau)\big) + \sum_{\tau \not\equiv k \,(\mathrm{mod}\, p)} a_*(\tau) \\ &= n^2 - \sum_{\tau \equiv 0 \,(\mathrm{mod}\, p)} \big(a(\tau) - a_*(\tau)\big) - \sum_{\tau \equiv k \,(\mathrm{mod}\, p)} a_*(\tau), \end{aligned} \tag{2.11}$$

where $p = \mathrm{char}\, \mathbb{F}_q$. Therefore, $C(n,k;q)$ is determined by $a(\tau)$ and $a_*(\tau)$, $0 \le \tau \le k$.

Proposition 2.6. (i) *Let $0 \le i \le n-1$. We have*

$$a_i(\tau) = \frac{1}{\tau} \big|\{x \in K_i \setminus \{-1\} : |S^{-1}(x)| = \tau - 1\}\big|, \quad 2 \le \tau \le k, \tag{2.12}$$

$$a_i(1) = k - \delta_i - \sum_{\tau=2}^{k} \tau a_i(\tau), \tag{2.13}$$

$$a_i(0) = n - \sum_{\tau=1}^{k} a_i(\tau). \tag{2.14}$$

(ii) *We have*

$$a(\tau) = \frac{1}{\tau}\big|\{x \in \mathbb{Z}_r^* \setminus \{-1\} : |S^{-1}(x)| = \tau - 1\}\big|, \quad 2 \le \tau \le k, \tag{2.15}$$

$$a(1) = nk - 1 - \sum_{\tau=2}^{k} \tau a(\tau), \tag{2.16}$$

$$a(0) = n^2 - \sum_{\tau=1}^{k} a(\tau). \tag{2.17}$$

Proof. (i) We only have to prove (2.12). ((2.13) follows from (2.8), and (2.14) is obvious.) Let $2 \le \tau \le k$ and let

$$\mathcal{X} = \{x \in K_i \setminus \{-1\} : |S^{-1}(x)| = \tau - 1\}.$$

Define

$$\begin{array}{rccl} f: & \mathcal{X} & \longrightarrow & \{0 \le j \le n-1 : t_{ij} = \tau\} \\ & x & \longmapsto & j, \quad \text{where } 1 + x \in K_j. \end{array}$$

1° We claim that f is well defined. Assume that $x \in \mathcal{X}$. Since $S^{-1}(x) \ne \emptyset$, $x \ne -1$, so $1 + x \in K_j$ for some $0 \le j \le n-1$. Note that

$$\begin{aligned} t_{ij} &= |(1 + xK) \cap (1 + x)K| \\ &= |[x]| && \text{(by (2.2))} \\ &= 1 + |S^{-1}(x)| && \text{(by (2.5))} \\ &= \tau. \end{aligned}$$

2° We claim that f is onto. Assume that $t_{ij} = \tau$. Since $\tau \ge 2$, by Lemma 2.1, there exists $x \in \mathbb{Z}_r^* \setminus \{-1\}$ such that $K_i = xK$ and $K_j = (1 + x)K$. Moreover, by (2.5) and (2.2),

$$|S^{-1}(x)| = |[x]| - 1 = |((1 + xK) \cap (1 + x)K| - 1 = t_{ij} - 1 = \tau - 1.$$

Hence $x \in \mathcal{X}$ and $f(x) = j$.

3° Let $x \in \mathcal{X}$ and $j = f(x)$. We claim that $f^{-1}(j) = [x]$; since $|[x]| = \tau$, this claim implies that f is τ-to-1. If $x' \in f^{-1}(j)$, then $x' \in K_i$ and $1 + x' \in K_j = (1 + x)K$, i.e., $x' \in [x]$. On the other hand, if $x' \in [x]$, then

$$|S^{-1}(x')| = |[x']| - 1 = |[x]| - 1 = \tau - 1.$$

Hence $x' \in \mathcal{X}$. Since $1 + x' \in (1 + x)K = K_j$, we have $f(x') = j$, i.e., $x \in f^{-1}(j)$.

4° Equation (2.12) follows from 1° – 3°.

(ii) The equations follow from (i). □

Assume that $t_{ij} \le 2$ for all $0 \le i, j \le n-1$, i.e., $S : \mathcal{S}_k \to \mathbb{Z}_r^* \setminus \{-1\}$ is one-to-one. Under this assumption, we are able to determine $a(2)$ and $a_*(2)$ explicitly. By (2.15),

$$a(2) = \frac{1}{2}|\mathcal{S}_k| = \frac{1}{2}(k-1)(k-2). \tag{2.18}$$

We claim that $1 \notin S(\mathcal{S}_k)$. If, to the contrary, $S(u,v) = 1$ for some $(u,v) \in \mathcal{S}_k$, then $S(u,v) = S(u,v)^{-1} = S(-u, v-u)$, where $(-u, v-u) \in \mathcal{S}_k$. It follows that $(u,v) = (-u, v-u)$, which is a contradiction.

(i) Assume that k is even. Then $\{\omega^0, \omega^{k/2}\} = \{\pm 1\}$ is disjoint from $S(\mathcal{S}_k)$. On the other hand, for all $u \in \mathbb{Z}_k \setminus \{0, k/2\}$, $\omega^u = S(-2u, -u)$, where $(-2u, -u) \in \mathcal{S}_k$. Hence

$$a_0(2) = \frac{1}{2}(k-2). \tag{2.19}$$

(ii) Assume that k is odd. We claim that $S(\mathcal{S}_k) \cap (-K) = \emptyset$. If, to the contrary, there exist $(u,v) \in \mathcal{S}_k$ and $l \in \mathbb{Z}_k$ such that $S(u,v) = -\omega^l$. It follows that

$$-\frac{1-\omega^v}{\omega^{2v-u} - \omega^v} = \omega^{l+u-v} = -\frac{1-\omega^{-(l+u-v)}}{\omega^{-2(l+u-v)} - \omega^{-(l+u-v)}}.$$

In the above, $\omega^{l+u-v} \neq -1$ since k is odd. Thus $2v - u \neq 0$ and $(2v-u, v) \in \mathcal{S}_k$. Moreover, $l + u - v \neq 0$ since otherwise, $S(2v-u, v) = 1$, which is impossible. Therefore $(-2(l+u-v), -(l+u-v)) \in \mathcal{S}_k$ and

$$S(2v-u, u) = S(-2(l+u-v), -(l+u-v)).$$

Then $(2v-u, v) = (-2(l+u-v), -(l+u-v))$, which forces $u = 0$, a contradiction. Therefore the claim is proved. We conclude that

$$a_{n/2}(2) = 0. \tag{2.20}$$

Remark 2.7. In the above, one can further determine $a(\tau)$ and $a_*(\tau)$, $\tau = 0, 1$, using Proposition 2.6. Then $C(n,k;q)$ is given by (2.11), and the result is Theorem 1.1.

3. Exceptional Primes

We follow the notation of Section 2. First note that for $(u,v), (u'v') \in \mathcal{S}_k$, $S(u,v) = S(u',v')$ if and only if $\overline{f}_{(u,v),(u'v')}(\omega) = 0$, where

$$\begin{aligned} f_{(u,v),(u'v')} &= X^{u+v'} - X^{u'+v} - X^u + X^{u'} + X^v - X^{v'} \\ &= (1-X^u)(1-X^{v'}) - (1-X^{u'})(1-X^v) \in \mathbb{Z}[X], \end{aligned} \tag{3.1}$$

and $\overline{(\ )}$ is the reduction from $\mathbb{Z}[X]$ to $\mathbb{Z}_r[X]$. Let $\Phi_k \in \mathbb{Z}[X]$ denote the kth cyclotomic polynomial. Then

$$\overline{\Phi}_k = \prod_{l \in \mathbb{Z}_k^\times} (X - \omega^l),$$

where $\mathbb{Z}_k^\times$ is the multiplicative group of $\mathbb{Z}_k$.

Lemma 3.1. *Let $(u,v), (u',v') \in \mathcal{S}_k$ be such that $(u,v) \neq (u',v')$. Then*

$$f_{(u,v),(u'v')} \not\equiv 0 \pmod{\Phi_k}.$$

Proof. Assume to the contrary that $f_{(u,v),(u'v')} \equiv 0 \pmod{\Phi_k}$. Then

$$(\zeta_k^u - 1)(\zeta_k^{v'} - 1) = (\zeta_k^{u'} - 1)(\zeta_k^v - 1), \tag{3.2}$$



FIGURE 1. Polar decomposition of $\zeta_k^u - 1$

where $\zeta_k = e^{2\pi i/k}$. We treat u, v, u', v' as elements of $\{1, 2, \ldots, k-1\}$. The polar decomposition of $\zeta_k^u - 1$ is

$$\zeta_k^u - 1 = 2\sin\frac{\pi u}{k} \cdot e^{i(\frac{\pi u}{k} + \frac{\pi}{2})};$$

see Figure 1. Then (3.2) is equivalent to the following system:

$$\sin\frac{\pi u}{k}\sin\frac{\pi v'}{k} = \sin\frac{\pi u'}{k}\sin\frac{\pi v}{k}, \tag{3.3}$$

$$\frac{\pi u}{k} + \frac{\pi v'}{k} = \frac{\pi u'}{k} + \frac{\pi v}{k}. \tag{3.4}$$

By (3.3),

$$\cos\frac{\pi}{k}(u - v') - \cos\frac{\pi}{k}(u + v') = \cos\frac{\pi}{k}(u' - v) - \cos\frac{\pi}{k}(u' + v).$$

Thus by (3.4),

$$\cos\frac{\pi}{k}(u - v') = \cos\frac{\pi}{k}(u' - v),$$

i.e.,

$$u - v' = \pm(u' - v). \tag{3.5}$$

Combining (3.4) and (3.5) gives

$$\begin{cases} u + v' = u' + v, \\ u - v' = \pm(u' - v). \end{cases}$$

It follows that either $(u, v) = (u', v')$ or $(u, u') = (v, v')$, neither of which is possible. □

Let $\mathcal{P}_k$ denote the set of primes $r > k+1$ such that $r \equiv 1 \pmod{k}$ and $S : \mathcal{S}_k \to \mathbb{Z}_r^* \setminus \{-1\}$ is not one-to-one. Recall that the elements of $\mathcal{P}_k$ are called exceptional primes. Moreover, for each $m \in \mathbb{Z}$, let $\mathcal{P}_k(m)$ be the set of primes divisors r of m such that $r > k+1$ and $r \equiv 1 \pmod{k}$.

Theorem 3.2. *We have*

$$\mathcal{P}_k = \mathcal{P}_k\Bigg(\prod_{\substack{(u,v),(u',v') \in \mathcal{S}_k \\ (u,v) \neq (u',v')}} \mathrm{Res}(f_{(u,v),(u',v')}, \Phi_k)\Bigg), \tag{3.6}$$

where $\mathrm{Res}(\cdot\,,\cdot)$ *is the resultant.*

Remark 3.3. By Lemma 3.1, the product in (3.6) is a nonzero integer. Hence it follows from (3.6) that $|\mathcal{P}_k| < \infty$.

Proof of Theorem 3.2. Let $\mathcal{R}$ denote the right side of (3.6).

First assume that $r \in \mathcal{P}_k$. Then there exist $(u,v),(u',v') \in \mathcal{S}_k$ such that $(u,v) \neq (u',v')$ and $\overline{f}_{(u,v),(u',v')}(\omega) = 0$, where $\overline{(\ )}$ is the reduction from $\mathbb{Z}[X]$ to $\mathbb{Z}_r[X]$. Then

$$\overline{\mathrm{Res}(f_{(u,v),(u',v')}, \Phi_k)} = \mathrm{Res}(\overline{f}_{(u,v),(u',v')}, \overline{\Phi}_k) = 0,$$

and hence $r \in \mathcal{R}$.

Next assume that $r \in \mathcal{R}$. Then there exist $(u,v),(u',v') \in \mathcal{S}_k$ such that $(u,v) \neq (u',v')$ and $\mathrm{Res}(\overline{f}_{(u,v),(u',v')}, \overline{\Phi}_k) = 0$. It follows that $\overline{f}_{(u,v),(u',v')}(\omega^l) = 0$ for some $l \in \mathbb{Z}_k^\times$. Then $\overline{f}_{(lu,lv),(lu',lv')}(\omega) = 0$, where $(lu,lv),(lu',lv') \in \mathcal{S}_k$ and $(lu,lv) \neq (lu',lv')$. □

Since $f_{(u,v),(u',v')} = -f_{(u',v'),(u,v)} = -f_{(v,u),(v',u')}$, for the product in (3.6), we only have to consider those $(u,v),(u',v') \in \mathcal{S}_k$ such that $u \leq \min\{v,u',v'\}$. Moreover, we may further exclude those $(u,v),(u',v')$ with $u = u'$ or $v = v'$. In fact, if say $v = v'$, then for any prime $r \equiv 1 \pmod{k}$, $\overline{f}_{(u,v),(u',v')}(\omega^l) = (\omega^{lu'} - \omega^{lu})(1-\omega^{lv})$ is never 0 for all $l \in \mathbb{Z}_k^\times$. Hence

$$\mathrm{Res}(f_{(u,v),(u',v')}, \Phi_k) \not\equiv 0 \pmod{r}. \tag{3.7}$$

If one prefers a more direct argument for (3.7), observe that

$$\begin{aligned} &\mathrm{Res}(f_{(u,v),(u',v')}, \Phi_k) \\ &= \mathrm{Res}\big(-X^u(X^{u'-u}-1)(X^v-1), \Phi_k\big) \\ &= \pm \prod_{l \in \mathbb{Z}_k^\times} (\zeta_k^{(u'-u)l} - 1)(\zeta_k^{vl} - 1) \\ &= \pm\, \Phi_{k/\gcd(k,u'-u)}(1)^{\gcd(k,u'-u)} \Phi_{k/\gcd(k,v)}(1)^{\gcd(k,v)} \end{aligned} \tag{3.8}$$

For each integer $m > 1$, $\Phi_m \mid 1 + X + \cdots + X^{m-1}$ and $\Phi_m(1) \mid m$. Therefore by (3.8), $\mathrm{Res}(f_{(u,v),(u',v')}, \Phi_k)$ does not have any prime divisor $> k$.

We conclude that

$$\mathcal{P}_k = \mathcal{P}_k \Bigg(\prod_{\substack{(u,v),(u',v') \in \mathcal{S}_k \\ v,u' > u,\ v' \geq u,\ v \neq v'}} \mathrm{Res}(f_{(u,v),(u',v')}, \Phi_k) \Bigg). \tag{3.9}$$

When $k \leq 3$, $|\mathcal{S}_k| \leq 1$, so the product in (3.9) is an empty one and hence $\mathcal{P}_k = \emptyset$. The sets $\mathcal{P}_k$, $4 \leq k \leq 20$, are given in Table 1 in the next section.

4. Computation of $C(n,k;q)$ for Exceptional Primes

Algorithm 4.1. Given an integer $k > 0$ and an exceptional prime $r = nk+1 \in \mathcal{P}_k$, the following steps produce the complexity $C(n,k;q)$.

Step 1. Find $\omega \in \mathbb{Z}_r^*$ with $o(\omega) = k$. Compute the multiset $\mathfrak{S} = \{S(u,v) : (u,v) \in \mathcal{S}_k\}$. Each element of $\mathfrak{S}$ is an integer in $\{1,2,\ldots,r-2\}$.

Step 2. Tally the multiset $\mathfrak{S}$ to determine the multiset $\mathfrak{M}$ of the multiplicities of the elements of $\mathfrak{S}$ and the multiset $\mathfrak{M}_*$ of the multiplicities of the elements in $\mathfrak{S}$ that belong to $(-1)^k\langle\omega\rangle$. More precisely, if

$$\mathfrak{S} = \{\underbrace{x_1,\dots,x_1}_{m_1},\dots,\underbrace{x_a,\dots,x_a}_{m_a},\dots,\underbrace{x_b,\dots,x_b}_{m_b}\},$$

where $x_1,\dots,x_b$ are distinct and $\{x_1,\dots,x_b\}\cap(-1)^k\langle\omega\rangle = \{x_1,\dots,x_a\}$, then $\mathfrak{M} = \{m_1,\dots,m_b\}$ and $\mathfrak{M}_* = \{m_1,\dots,m_a\}$.

Step 3. For each $2 \le \tau \le k$, compute

$$a(\tau) = \frac{1}{\tau}\cdot(\text{the multiplicity of } \tau-1 \text{ in } \mathfrak{M}), \tag{4.1}$$

$$a_*(\tau) = \frac{1}{\tau}\cdot(\text{the multiplicity of } \tau-1 \text{ in } \mathfrak{M}_*). \tag{4.2}$$

Also compute

$$a(1) = nk-1-\sum_{\tau=2}^{k}\tau a(\tau), \qquad a(0) = n^2-\sum_{\tau=1}^{k}a(\tau), \tag{4.3}$$

$$a_*(1) = k-1-\sum_{\tau=2}^{k}\tau a_*(\tau), \qquad a_*(0) = n-\sum_{\tau=1}^{k}a_*(\tau). \tag{4.4}$$

Step 4. Compute

$$C(n,k;q) = n^2 - \sum_{\tau\equiv 0 \,(\mathrm{mod}\, p)}\big(a(\tau)-a_*(\tau)\big) - \sum_{\tau\equiv k \,(\mathrm{mod}\, p)} a_*(\tau). \tag{4.5}$$

Note. In the above, (4.1) – (4.4) are from Proposition 2.6, and (4.5) is (2.11). The algorithm works for all primes $r = nk+1$, but it is not necessary if $r \notin \mathcal{P}_k$ because of Theorem 1.1.

The above algorithm is applied to $4 \le k \le 20$ and $r \in \mathcal{P}_k$. The sequences $a(\tau)$ and $a_*(\tau)$ are given in Table 1. Step 4 is easy computation but the results are too lengthy to be included. Instead, we use examples to exhibit the formulas for $C(n,k;q)$ for a few pairs of parameters (n,k).

Example 4.2. Let $k=6$, $n=3$, $r=19$. The nonzero terms of the sequences $a(\tau)$ and $a_*(\tau)$ are

$$a(1)=3, \quad a(2)=4, \quad a(3)=2,$$
$$a_*(1)=1, \quad a_*(2)=2.$$

By (4.5),

$$C(3,6;q) = \begin{cases} 5 & \text{if } p=2,\\ 7 & \text{if } p=3,\\ 8 & \text{if } p=5,\\ 9 & \text{if } p>5. \end{cases} \tag{4.6}$$

Note that (4.6) is different from the result in [3, Theorem 3.14]. It appears that the exceptional prime $r=19$ for $k=6$ was not detected in [3].

Example 4.3. Let $k = 7$, $n = 6$, $r = 43$. The nonzero terms of the sequences $a(\tau)$ and $a_*(\tau)$ are

$$a(0) = 9, \quad a(1) = 14, \quad a(2) = 12, \quad a(3) = 1,$$
$$a_*(0) = 2, \quad a_*(1) = 2, \quad a_*(2) = 2.$$

By (4.5),

$$C(6, 7; q) = \begin{cases} 26 & \text{if } p = 3, \\ 27 & \text{if } p = 5, 7, \\ 29 & \text{if } p > 7. \end{cases} \tag{4.7}$$

The prime $p = 2$ is not included in (4.7) since the order of 2 in $\mathbb{Z}_{43}^*$ is 14 and $\gcd(nk/14, n) \neq 1$. It appears that the exceptional prime $r = 43$ for $k = 7$ was not detected in [8, Theorem 10].

Example 4.4. Let $k = 20$, $n = 1166$, $r = 23321$. The nonzero terms of the sequences $a(\tau)$ and $a_*(\tau)$ are

$$a(0) = 1336402, \quad a(1) = 22995, \quad a(2) = 153, \quad a(3) = 6,$$
$$a_*(0) = 1156, \quad a_*(1) = 1, \quad a_*(2) = 9.$$

By (4.5),

$$C(1166, 20; q) = \begin{cases} 24295 & \text{if } p = 3, \\ 24310 & \text{if } p = 17 \text{ or } p > 19. \end{cases} \tag{4.8}$$

The primes $p = 2, 5, 7, 11, 13, 19$ are not included in (4.8) since $\gcd(nk/e(p, r), n) \neq 1$ for these primes p, where $e(p, r)$ is the order of p in $\mathbb{Z}_r^*$.

Table 1 suggests that for every prime $r = 2k + 1$, $r \in \mathcal{P}_k$. Moreover, in this case, the nonzero terms of $a(\tau)$ and $a_*(\tau)$ are

$$\begin{cases} a(k/2 - 1) = 1, \quad a(k/2) = 3, \\ a_*(k/2 - 1) = 1, \quad a_*(k/2) = 1 \end{cases} \qquad \text{if } k \text{ is even}, \tag{4.9}$$

$$\begin{cases} a((k-1)/2) = 3, \quad a((k+1)/2) = 1, \\ a_*((k-1)/2) = 2 \end{cases} \qquad \text{if } k \text{ is odd}. \tag{4.10}$$

The above are indeed correct formulas and they follow from [6, §6]. The following is a proof using the notation of the present paper.

Let $(u, v) \in \mathcal{S}_k$ and $z \in \mathbb{Z}_r \setminus \{0, -1\}$. Write $\omega^u = x^2$ and $\omega^v = y^2$, where $x, y \in \mathbb{Z}_r \setminus \{0, \pm 1\}$. Then $S(u, v) = z$ if and only if

$$\frac{x^2 - 1}{y^2 - 1} = 1 + z^{-1},$$

i.e.,

$$x^2 - \lambda y^2 = 1 - \lambda, \tag{4.11}$$

where $\lambda = 1 + z^{-1} \in \mathbb{Z}_r \setminus \{0, 1\}$. By [9, Theorem 6.26],

$$|\{(x, y) \in \mathbb{Z}_r^2 : x^2 - \lambda y^2 = 1 - \lambda\}| = r - \eta(\lambda),$$

where η is the quadratic character of $\mathbb{Z}_r$. It follows that

$$\left|\left\{(x, y) : x, y \in \mathbb{Z}_r \setminus \{0, \pm 1\},\ x^2 - \lambda y^2 = 1 - \lambda\right\}\right|$$

$$\begin{aligned} &= r - \eta(\lambda) - 4 - (1 + \eta(1-\lambda)) - (1 + \eta(1-\lambda^{-1})) \\ &= r - 6 - \eta(\lambda) - \eta(1-\lambda) - \eta(1-\lambda^{-1}). \end{aligned} \tag{4.12}$$

First assume that k is even. Then $\eta(\lambda) + \eta(1-\lambda) + \eta(1-\lambda^{-1}) = -1$ or 3 since the left side of (4.12) is $\equiv 0 \pmod 4$. Let

$$b_i = \big|\big\{\lambda \in \mathbb{Z}_r \setminus \{0,1\} : \eta(\lambda) + \eta(1-\lambda) + \eta(1-\lambda^{-1}) = i\big\}\big|, \quad i = -1, 3.$$

Then

$$b_{-1} + b_3 = r - 2 \tag{4.13}$$

and

$$-b_{-1} + 3b_3 = \sum_{\lambda \in \mathbb{Z}_r \setminus \{0,1\}} \big(\eta(\lambda) + \eta(1-\lambda) + \eta(1-\lambda^{-1})\big) = -3. \tag{4.14}$$

Combining (4.13) and (4.14) gives

$$b_{-1} = \frac{3}{4}(r-1), \quad b_3 = \frac{1}{4}(r-5).$$

For $i = -1, 3$, by (4.12),

$$\left|\left\{z \in \mathbb{Z}_r \setminus \{0,-1\} : |S^{-1}(z)| = \frac{1}{4}(r-6-i)\right\}\right| = b_i.$$

Hence

$$a((r-6-i)/4+1) = \frac{1}{(r-6-i)/4+1} b_i,$$

i.e., $a(k/2-1)+1) = 1$ and $a(k/2) = 3$.

To determine $a_*((r-6-i)/4+1)$ with $i = -1, 3$, let

$$\begin{aligned} b_i^* &= \big|\big\{\lambda \in \mathbb{Z}_r \setminus \{0,1\} : \eta(\lambda-1) = 1,\ \eta(\lambda) + \eta(1-\lambda) + \eta(1-\lambda^{-1}) = i\big\}\big| \\ &= \big|\big\{\lambda \in \mathbb{Z}_r \setminus \{0,1\} : \eta(\lambda-1) = 1,\ \eta(\lambda) = (i-1)/2\big\}\big|. \end{aligned}$$

In the above, let $1 - \lambda = x^2$ and $\lambda = \epsilon y^2$, where $x, y \in \mathbb{Z}_r^*$ and $\epsilon \in \mathbb{Z}$ is a fixed element such that $\eta(\epsilon) = (i-1)/2$. Then

$$\begin{aligned} b_i^* &= \frac{1}{4}|\{(x,y) \in \mathbb{Z}_r^2 : x^2 \neq 0, 1,\ x^2 + \epsilon y^2 = 1\}| \\ &= \frac{1}{4}\big(r - \eta(-\epsilon) - (1 + \eta(\epsilon)) - 2\big) \qquad \text{(by [9, Theorem 6.26])} \\ &= \frac{1}{4}(r - 3 - 2\eta(\epsilon)) \\ &= \frac{1}{4}(r - 2 - i). \end{aligned}$$

Thus

$$a_*((r-6-i)/4+1) = \frac{1}{(r-6-i)/4+1} b_i^* = 1,$$

i.e., $a_*(k/2-1) = a_*(k/2) = 1$.

Next assume that k is odd. Then in (4.12), $\eta(\lambda) + \eta(1-\lambda) + \eta(1-\lambda^{-1}) = -3$ or 1. Let

$$c_i = \big|\big\{\lambda \in \mathbb{Z}_r \setminus \{0,1\} : \eta(\lambda) + \eta(1-\lambda) + \eta(1-\lambda^{-1}) = i\big\}\big|, \quad i = -3, 1.$$

Then

$$\begin{cases} c_{-3} + c_1 = r - 2, \\ -3c_{-3} + c_1 = -3, \end{cases}$$

which gives

$$c_{-3} = \frac{1}{4}(r+1), \quad c_1 = \frac{3}{4}(r-3).$$

We have

$$a((r-6-i)/4+1) = \frac{1}{(r-6-i)/4+1} c_i,$$

i.e., $a((k-1)/2) = 3$, $a((k+1)/2) = 1$. To determine $a_*((r-6-i)/4+1)$ with $i = -3, 1$, let

$$c_i^* = \left|\left\{\lambda \in \mathbb{Z}_r \setminus \{0,1\} : \eta(\lambda - 1) = -1,\ \eta(\lambda) + \eta(1-\lambda) + \eta(1-\lambda^{-1}) = i\right\}\right|.$$

Note that $\eta(\lambda - 1) = -1$ implies that $\eta(\lambda) + \eta(1-\lambda) + \eta(1-\lambda^{-1}) = 1$. Hence

$$c_i^* = \begin{cases} 0 & \text{if } i = -3, \\ |\{\lambda \in \mathbb{Z}_r \setminus \{0,1\} : \eta(\lambda - 1) = -1\}| = \dfrac{r-3}{2} & \text{if } i = 1. \end{cases}$$

Thus

$$a_*((r-6-i)/4+1) = \frac{1}{(r-6-i)/4+1} c_i^*,$$

i.e., $a_*((k-1)/2) = 2$ and $a_*((k+1)/2) = 0$.

References

[1] V. V. Acharya and S. A. Katre, *Cyclotomic numbers of orders 2l, l an odd prime*, Acta Arith., **69** (1995), 51 – 74.

[2] L. D. Baumert and H. Fredricksen, *The cyclotomic number of order eighteen with applications to difference sets*, Math. Comp., **21** (1967), 204 – 219.

[3] M. Christopoulou, T. Garefalakis, D. Panario, D. Thomson, *Gauss periods as constructions of low complexity normal bases*, Des. Codes Cryptogr., **62** (2012), 43 – 62.

[4] S. Gao, *Normal Bases over Finite Fields*, Ph.D. Dissertation, University of Waterloo, 1993.

[5] S. Gao, J. von zur Gathen, D. Panario, V. Shoup, *Algorithms for exponentiation in finite fields*, J. Symbolic Computation, **29** (2000), 879 – 889.

[6] L. E. Dickson, *Cyclotomy, higher Congruences, and Waring's Problem*, Amer. J. Math., **57** (1935), 391 – 424.

[7] S. A. Katre and A. R. Rajwade, *Complete solution of the cyclotomic problem in $\mathbb{F}_q$ for any prime modulus l, $q = p^\alpha$, $p \equiv 1 \pmod{l}$*, Acta Arith., **45** (1985), 183 – 199.

[8] Q. Y. Liao and X. L. Hu, *An explicit formula for the complexity of a class of Gauss period normal bases over finite fields* (in Chinese), Acta Mathematica Sinica, Chinese Series, **57** (2014), 863 – 874.

[9] R. Lidl and H. Niederreiter, *Finite Fields*, Cambridge Univ. Press, New York, 1997.

[10] R. C. Mullin, I. M. Onyszchuk, S. A. Vanstone, R. M. Wilson, *Optimal normal bases in* GF(p^n), Discrete Applied Math., **22** (1988/89), 149 – 161.

[11] J. B. Muskat, *The cyclotomic numbers of order fourteen*, Acta Arith., **11** (1965/1966), 263 – 279.

[12] D. Shirolkar and S. A. Katre, *Jacobi sums and cyclotomic numbers of order l^2*, Acta Arith., **147** (2011), 33 – 49.

[13] A. Wassermann, *Konstruktion von Normalbasen*, Bayreuther Mathematische Schriften, **31** (1990), 155 – 164.

[14] A. L. Whiteman, *The cyclotomic numbers of order twelve*, Acta Arith., **6** (1960), 53 – 76.

Department of Mathematics and Statistics, University of South Florida, Tampa, FL 33620

E-mail address: xhou@usf.edu

TABLE 1. Values of $a(\tau)$ and $a_*(\tau)$ for $4 \le k \le 20$, $r = nk+1 \in \mathcal{P}_k$

The top sequence is $a(\tau)$, and the bottom one is $a_*(\tau)$, $0 \le \tau \le 10$; only nonzero terms are listed

k	n	0	1	2	3	4	5	6	7	8	9	10
4	–											
5	2			3	1							
				2								
6	2			1	3							
				1	1							
6	3		3	4	2							
			1	2								
7	4	1	6	6	3							
			2	2								
7	6	9	14	12	1							
		2	2	2								
8	2				1	3						
					1	1						
8	5	4	6	12	3							
		2		2	1							
9	2					3	1					
						2						
9	4		3	10		3						
				4								
9	8	15	33	10	6							
			8									
9	12	62	60	19	3							
		6	4	2								
9	14	96	78	19	3							
		8	4	2								
9	30	657	219	22	2							
		22	8									
10	3			3	1	5						
				1	1	1						
10	4	1		9	3	3						
		1		1	1	1						
10	6	4	12	15	3	2						
		2		3	1							
10	7	10	15	18	6							
		2	1	4								
10	10	31	45	18	6							
		5	1	4								
11	2						3	1				
							2					
11	6	1	15	15		5						
			2	4								
11	8	13	21	27		3						
		2	2	4								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
11	18	169	116	36	3							
		10	6	2								
11	32	712	279	27	6							
		22	10									
11	36	943	314	36	3							
		28	6	2								
11	62	3207	594	42	1							
		54	6	2								
12	3			1	2	3	3					
				1		1	1					
12	5	3		10	9	3						
		1		2	1	1						
12	6	5	6	10	15							
		1		4	1							
12	8	12	21	19	12							
		2	1	5								
12	9	23	15	37	6							
		3	1	5								
12	13	63	63	37	6							
		7	1	5								
12	15	99	75	49	2							
		9	1	5								
12	16	114	99	37	6							
		10	1	5								
13	4			6	3	6		1				
				2		2						
13	6	1	11	12	6	6						
			2	2	2							
13	10	22	39	30	6	3						
		2	4	4								
13	12	40	63	36		5						
		4	4	4								
13	24	322	203	48		3						
		16	4	4								
13	40	1144	396	57	3							
		30	8	2								
13	46	1579	483	48	6							
		34	12									
13	66	3559	743	48	6							
		54	12									
13	100	8761	1185	48	6							
		88	12									
13	124	13828	1488	57	3							
		114	8	2								
13	154	21778	1878	57	3							
		144	8	2								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
13	210	41436	2600	63	1							
		200	8	2								
14	2							1	3			
								1	1			
14	3				1	5		3				
					1	1		1				
14	5	1		12	6	3	3					
		1		2		1	1					
14	8	7	24	15	15	3						
		3		3	1	1						
14	9	18	14	39	7	3						
		4		3	1	1						
14	14	67	75	42	12							
		7	1	6								
14	15	82	87	48	6	2						
		8	1	6								
14	17	118	114	51	3	3						
		12		3	1	1						
14	20	187	159	42	12							
		13	1	6								
14	24	313	197	60	6							
		17	1	6								
14	27	424	239	60	6							
		20	1	6								
14	35	808	351	60	6							
		28	1	6								
14	39	1045	414	57	3	2						
		33		5	1							
14	47	1624	519	60	6							
		40	1	6								
14	50	1873	561	60	6							
		43	1	6								
14	75	4648	911	60	6							
		68	1	6								
14	78	5068	944	69	3							
		72		5	1							
14	92	7249	1149	60	6							
		85	1	6								
15	2								3	1		
									2			
15	4			1	9	3		3				
					2	2						
15	10	21	27	37	12	3						
			6	4								
15	12	38	48	46	9	3						
		4	2	6								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
15	14	54	93	37	6	6						
		6	4	2	2							
15	16	93	99	55	6	3						
		6	6	4								
15	18	125	147	37	12	3						
		8	6	4								
15	22	234	180	63	6		1					
		14	4	2	2							
15	28	453	246	82	3							
		16	10	2								
15	36	833	399	55	6	3						
		26	6	4								
15	38	954	423	55	12							
		24	14									
15	42	1214	483	55	12							
		28	14									
15	50	1839	576	82	3							
		38	10	2								
15	88	6513	1146	82	3							
		76	10	2								
15	92	7173	1206	82	3							
		80	10	2								
15	98	8214	1323	55	12							
		84	14									
15	144	18662	1995	73	6							
		130	14									
15	150	20336	2085	73	6							
		136	14									
15	154	21492	2145	73	6							
		140	14									
15	186	31892	2625	73	6							
		172	14									
15	208	40230	2955	73	6							
		194	14									
15	242	55020	3465	73	6							
		228	14									
15	270	68940	3873	85	2							
		256	14									
15	380	138786	5535	73	6							
		366	14									
15	388	144810	5655	73	6							
		374	14									
15	392	147870	5715	73	6							
		378	14									
15	458	202980	6705	73	6							
		444	14									

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
15	3084	9464886	46083	85	2							
		3070	14									
16	6	1	9	3	12	11						
		1	1	1		3						
16	7	4	6	21	12	3	3					
		2		3		1	1					
16	12	37	44	42	21							
		5		6	1							
16	15	79	65	69	12							
		7	1	7								
16	16	90	90	66	7	3						
		10		4	1	1						
16	21	193	176	60	9	3						
		15		4	1	1						
16	22	226	177	69	12							
		14	1	7								
16	25	325	207	87	6							
		17	1	7								
16	27	397	239	87	6							
		19	1	7								
16	28	436	255	87	6							
		20	1	7								
16	36	820	383	87	6							
		28	1	7								
16	37	877	399	87	6							
		29	1	7								
16	40	1060	447	87	6							
		32	1	7								
16	55	2245	687	87	6							
		47	1	7								
16	76	4660	1023	87	6							
		68	1	7								
16	132	15412	1919	87	6							
		124	1	7								
16	133	15661	1935	87	6							
		125	1	7								
17	6		5	15		15		1				
				4		2						
17	8	7	12	21	15	9						
			4	2		2						
17	14	48	78	60	3	6		1				
		6	4	2		2						
17	18	118	125	66	12	3						
		6	8	4								
17	24	268	227	66	12	3						
		12	8	4								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
17	26	331	267	66	6	6						
		16	6	2	2							
17	36	790	413	84	6	3						
		24	8	4								
17	38	904	447	84	6	3						
		26	8	4								
17	54	2104	719	84	6	3						
		42	8	4								
17	56	2290	753	84	6	3						
		44	8	4								
17	60	2680	839	66	12	3						
		48	8	4								
17	80	5155	1137	102	6							
		64	16									
17	84	5734	1227	90		5						
		72	8	4								
17	98	8047	1461	84	12							
		82	16									
17	138	16807	2139	90	6	2						
		122	16									
17	168	25480	2639	102		3						
		156	8	4								
17	180	29455	2837	102	6							
		164	16									
17	194	34453	3075	102	6							
		178	16									
17	204	38263	3245	102	6							
		188	16									
17	210	40648	3338	111	3							
		196	12	2								
17	216	43093	3467	84	12							
		200	16									
17	344	112603	5625	102	6							
		328	16									
17	500	241615	8277	102	6							
		484	16									
17	546	288949	9059	102	6							
		530	16									
17	644	403903	10725	102	6							
		628	16									
17	708	489343	11813	102	6							
		692	16									
17	920	830878	15408	111	3							
		906	12	2								
17	1046	1076452	17550	111	3							
		1032	12	2								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
17	1140	1280335	19157	102	6							
		1124	16									
17	1484	2177146	24996	111	3							
		1470	12	2								
17	1548	2370106	26084	111	3							
		1534	12	2								
17	2054	4184113	34695	102	6							
		2038	16									
17	2570	6561330	43452	117	1							
		2556	12	2								
18	2									1	3	
										1	1	
18	4			3		3	7	3				
				1		1	1	1				
18	6	3		4	23	3		3				
		1		2	1	1		1				
18	7	3	6	16	12	9	3					
		1		4		1	1					
18	9	12	12	31	17	9						
		2		5	1	1						
18	10	15	27	28	24	6						
		1	1	8								
18	11	24	30	37	27	3						
		4		5	1	1						
18	15	72	57	76	20							
		6	1	8								
18	17	96	102	73	15	3						
		10		5	1	1						
18	21	182	159	82	18							
		12	1	8								
18	22	213	159	100	12							
		13	1	8								
18	24	269	195	100	12							
		15	1	8								
18	27	368	249	100	12							
		18	1	8								
18	29	444	285	100	12							
		20	1	8								
18	30	491	285	118	6							
		21	1	8								
18	32	579	321	118	6							
		23	1	8								
18	34	675	357	118	6							
		25	1	8								
18	41	1074	483	118	6							
		32	1	8								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
18	42	1142	492	127	3							
		34		7	1							
18	45	1346	555	118	6							
		36	1	8								
18	62	2859	861	118	6							
		53	1	8								
18	81	5234	1203	118	6							
		72	1	8								
18	121	12594	1923	118	6							
		112	1	8								
18	126	13743	2001	130	2							
		117	1	8								
18	127	13974	2031	118	6							
		118	1	8								
18	321	97398	5511	130	2							
		312	1	8								
19	10	15	15	48	12	9		1				
		2	2	4		2						
19	12	28	38	54	15	9						
		2	6	2		2						
19	22	193	189	81	18	3						
		8	10	4								
19	24	243	230	93	3	6		1				
		14	6	2		2						
19	30	460	329	99	6	6						
		18	8	2	2							
19	34	643	399	99	12	3						
		20	10	4								
19	40	979	495	117	6	3						
		26	10	4								
19	58	2392	861	99	6	6						
		46	8	2	2							
19	78	4741	1217	117	6	3						
		64	10	4								
19	82	5305	1293	117	6	3						
		68	10	4								
19	84	5593	1349	99	12	3						
		70	10	4								
19	94	7183	1539	99	12	3						
		80	10	4								
19	100	8239	1635	117	6	3						
		86	10	4								
19	108	9745	1805	99	12	3						
		94	10	4								
19	118	11821	1977	117	6	3						
		104	10	4								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
19	120	12259	2015	117	6	3						
		106	10	4								
19	142	17614	2409	135	6							
		124	18									
19	204	37882	3605	117	12							
		186	18									
19	232	49558	4137	117	12							
		214	18									
19	234	50449	4179	123		5						
		220	10	4								
19	240	53182	4289	117	12							
		222	18									
19	244	55048	4347	135	6							
		226	18									
19	258	61810	4613	135	6							
		240	18									
19	358	121504	6531	117	12							
		340	18									
19	360	122902	6567	123	6	2						
		342	18									
19	402	154108	7367	117	12							
		384	18									
19	460	203005	8457	135		3						
		446	10	4								
19	472	213964	8679	135	6							
		454	18									
19	492	232867	9050	144	3							
		476	14	2								
19	750	548392	13979	117	12							
		732	18									
19	778	590650	14493	135	6							
		760	18									
19	810	640858	15101	135	6							
		792	18									
19	852	709864	15899	135	6							
		834	18									
19	1620	2593768	30491	135	6							
		1602	18									
19	1678	2783950	31593	135	6							
		1660	18									
19	1834	3328858	34557	135	6							
		1816	18									
19	2112	4420564	39839	135	6							
		2094	18									
19	2128	4488100	40143	135	6							
		2110	18									

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
19	2532	6363064	47819	135	6							
		2514	18									
19	2824	7921471	53358	144	3							
		2808	14	2								
19	2892	8308864	54659	135	6							
		2874	18									
19	2970	8764621	56132	144	3							
		2954	14	2								
19	3352	11172364	63399	135	6							
		3334	18									
19	3880	14980828	73431	135	6							
		3862	18									
19	4402	19294117	83340	144	3							
		4386	14	2								
19	4852	23449864	91899	135	6							
		4834	18									
19	5188	26816920	98283	135	6							
		5170	18									
19	5638	31680073	106824	144	3							
		5622	14	2								
19	6888	47313823	130574	144	3							
		6872	14	2								
19	7372	54206464	139779	135	6							
		7354	18									
19	9198	84428595	174458	150	1							
		9182	14	2								
20	2										1	3
											1	1
20	3						3	1	2	3		
							1	1		1		
20	5	1			9	9		6				
		1			1	1		2				
20	9	4	15	39	12	9			2			
		2	1	3		3						
20	12	25	39	51	18	11						
		5	1	3		3						
20	14	43	63	63	18	9						
		7	1	3		3						
20	20	136	156	90	9	9						
		12		6	1	1						
20	21	163	167	81	30							
		11	1	9								
20	23	217	189	99	24							
		13	1	9								
20	26	310	231	117	18							
		16	1	9								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
20	27	328	284	105	6	3	3					
		20		5		1	1					
20	30	454	309	123	12	2						
		20	1	9								
20	32	538	351	117	18							
		22	1	9								
20	33	583	371	117	18							
		23	1	9								
20	35	679	411	117	18							
		25	1	9								
20	38	820	516	90	9	9						
		30		6	1	1						
20	41	1021	513	135	12							
		31	1	9								
20	44	1210	591	117	18							
		34	1	9								
20	47	1423	651	117	18							
		37	1	9								
20	51	1741	713	135	12							
		41	1	9								
20	53	1897	786	108	15	3						
		45		6	1	1						
20	59	2470	846	162	3							
		50		8	1							
20	60	2566	875	153	6							
		50	1	9								
20	65	3091	975	153	6							
		55	1	9								
20	66	3202	995	153	6							
		56	1	9								
20	68	3412	1086	108	15	3						
		60		6	1	1						
20	69	3541	1073	135	12							
		59	1	9								
20	74	4162	1155	153	6							
		64	1	9								
20	80	4954	1308	126	9	3						
		72		6	1	1						
20	81	5107	1295	153	6							
		71	1	9								
20	86	5842	1395	153	6							
		76	1	9								
20	87	5989	1433	135	12							
		77	1	9								
20	93	6943	1571	117	18							
		83	1	9								

TABLE 1. continued

k	n	0	1	2	3	4	5	6	7	8	9	10
20	95	7291	1575	153	6							
		85	1	9								
20	104	8902	1755	153	6							
		94	1	9								
20	122	12610	2115	153	6							
		112	1	9								
20	152	20230	2715	153	6							
		142	1	9								
20	156	21382	2795	153	6							
		146	1	9								
20	161	22867	2895	153	6							
		151	1	9								
20	173	26629	3153	135	12							
		163	1	9								
20	228	47590	4235	153	6							
		218	1	9								
20	251	58141	4713	135	12							
		241	1	9								
20	254	59602	4755	153	6							
		244	1	9								
20	311	90667	5895	153	6							
		301	1	9								
20	333	104395	6335	153	6							
		323	1	9								
20	348	114310	6635	153	6							
		338	1	9								
20	366	126802	6995	153	6							
		356	1	9								
20	408	158470	7835	153	6							
		398	1	9								
20	434	179842	8355	153	6							
		424	1	9								
20	686	457042	13395	153	6							
		676	1	9								
20	1166	1336402	22995	153	6							
		1156	1	9								